

## Original Research Article

## Determining the Effects of Ash from Hospital Waste Incineration on the Growth of *Fava Bean*

Khalid Mahmood Yosif <sup>1\*</sup><sup>1</sup>Biology Department College of Education for Pure Science University of Kirkuk, Kirkuk, Iraq**\*Corresponding Author:** Khalid Mahmood Yosif

Biology Department College of Education for Pure Science University of Kirkuk, Kirkuk, Iraq

**Article History**

Received: 16.01.2025

Accepted: 21.02.2025

Published: 24.02.2025

**Abstract:** The present study aimed to determine whether there are effects of medical waste incineration ash from hospitals in promoting the growth of fava bean plants. Samples were taken in soil (mixture) and ash in different proportions and placed in boxes according to sample numbers: 1 consisted of 1 kg of soil, sample number 2 also included 1 kg of ash, and sample number 3 contained half a kilo of soil plus half a kilo of ash. Sample number 4: a quarter of soil + 3 quarters of ash; sample number 5: a quarter of ash plus three quarters of soil. Before the research was done in the Directorate of Agriculture, Kirkuk, Department of Modern Irrigation Technologies, and Water and Soil Management Division Laboratories, a quantitative analysis of medical waste ash was done to find out the amounts of different materials and compounds in the ash. The analysis revealed a percentage of 0.189% for total nitrogen, 0.087% for total phosphorus, 0.47% for total potassium, and 25.476% for organic matter. We observed normal plant growth when we planted the seeds in soil, but no growth at all when we planted them in ash. When planted in an anvil containing half a kilo of soil and half a kilo of ash, the plant grew successfully, and no seeds were formed. In the fourth treatment, which included a quarter kilo of soil and three quarters of a kilo of ash, the plant did not grow, and on the contrary, it led to the appearance of white spots on the mixture. In the fifth treatment, which included a quarter kilo of ash and three quarters of a kilo of soil, the plant did not grow, and on the contrary, it led to the appearance of blackened soil. It is concluded from the current study that contaminated and hazardous waste should be buried in relation to medical waste, and restrictions should be placed on access to the waste disposal site.

**Keywords:** Hospital Waste, Ash, Bean, Nitrogen.

### INTRODUCTION

Medicinal waste encompasses all waste generated during diagnostic or medicinal procedures [1-3]. Medical waste, for instance, is defined as solid, liquid, and gaseous waste produced in hospitals, vet clinics, dentistry offices, medical research facilities, pharmaceutical companies, and blood banks [4, 5]. There are different types of medical waste, such as pathological waste, which includes human organs, body parts, and tissue; sharps, which includes needles, syringes, and blades; chemicals, which includes heavy metals from medical devices and solvents used in lab work; and infectious waste, which includes bodily fluids. Between 70 and 90 percent of medical waste is innocuous and similar to home waste. Hazardous categories classify the remaining 10–25%, raising concerns for human health and the environment [6, 7]. There are two primary categories of medical waste incinerator products. The waste that is discharged into the environment, including fly ash (MWFA), carbon dioxide, sulfur oxides, and chlorides, makes up the first section. The second component is bottom ash (MWBA), which is the ash that remains in the incinerator. The ash still in the incineration chamber makes up 75–90% of the total ash [8, 9]. We conducted several chemical investigations to study and assess the nature of the fly and bottom ash. The main studies found out about the ash's mineralogy, metal leaching, particle size distribution, and metal content [9]. Worldwide, there is no unified legislative framework governing the use of incineration ash. The laws governing the use of ash for fertilization vary from one nation to another. Lithuania only specifies the use of wood biomass ash. Certain chemical elements have minimum and maximum amounts in some nations. Finland, for instance, establishes minimum levels of calcium, phosphorus, and potassium in fertilizer made from biomass ash. Fly ash used in agriculture

**Copyright © 2025 The Author(s):** This is an open-access article distributed under the terms of the Creative Commons Attribution **4.0 International License (CC BY-NC 4.0)** which permits unrestricted use, distribution, and reproduction in any medium for non-commercial use provided the original author and source are credited.

**Citation:** Khalid Mahmood Yosif, (2025) Determining the Effects of Ash from Hospital Waste Incineration on the Growth of *Fava Bean*. *South Asian Res J Bio Appl Biosci*, 7(1), 46-50. 46

must have a minimum of 10% calcium content and a minimum of 2% total phosphorus and potassium content [10, 11]. Ash can be used in many different ways, depending on the type of biomass that is used. This is because biomass feedstocks are very different, and so are the ash that is made by burning them. The suitability of multi-crop biomass for solid biofuel production has already been supported by a number of research studies conducted by the authors [12-15]. Therefore, in the present study, the aim was to determine whether there are effects of ash from hospital waste incineration on the growth of fava bean plants.

## MATERIALS AND METHODS

When waste is burned in hospitals by the incinerator, it becomes ash after being isolated according to the nature of the material or compound used. The incinerator device must be handled with extreme caution, especially after burning the waste, and it requires a specialized cadre with a full scoop for burning waste. Medical waste refers to materials that have no value or use, yet their continued presence in the environment poses serious risks to human, animal, and plant sources and elements [16]. We took samples in different proportions of soil and ash, then placed them in a boxes.

Sample No. 1 / 1 kg of soil

Sample No. 2 / 1 kg of ash

Sample No. (3) / half a kilo of soil + half a kilo of ash

Sample No. (4) A quarter of soil + 3 quarters of ash

Sample No. (5) A quarter of ash + 3 quarters of soil

Then, the seeds of fava beans were used because they are considered important crops because their seeds are soft and dry. Most researchers were interested in this crop because it is considered an important food source for humans and sometimes animal feed. It also improves soil properties through nitrogen fixation and contributes to regulating the agricultural cycle, especially in areas that depend on rain (rain-fed areas).

## RESULTS AND DISCUSSION

Before the research was done in the Directorate of Agriculture, Kirkuk, Department of Modern Irrigation Technologies, and Water and Soil Management Division Laboratories, a quantitative analysis of medical waste ash was done to find out the amounts of different materials and compounds in the ash. Table 1 revealed that the percentage of total nitrogen was 0.189%, total phosphorus was 0.087%, total potassium was 0.47%, and organic matter was 25.476%.

We mixed the detected ash components with soil to see if any seed growth occurred after planting. It was found that when the seeds were planted in soil only (Figure 1), there was normal plant growth, while when planted with ash only (Figure 2), there was no growth. When planted in an anvil containing half a kilo of soil and half a kilo of ash (Figure 3), the plant grew successfully and no seeds were formed. In the fourth treatment, which included a quarter kilo of soil and three quarters of a kilo of ash (Figure 4), the plant did not grow, and on the contrary, it led to the appearance of white spots on the mixture. In the fifth treatment (Figure 5), which included a quarter kilo of ash and three quarters of a kilo of soil, the plant did not grow, and on the contrary, it led to the appearance of blackened soil.



**Figure 1: Plant growth normally when using only soil**



**Figure 2: Non-plant growth when using only ash**



**Figure 3: Plant growth when using half a kilo of soil and half a kilo of ash**



**Figure 4: non-plant growth when using quarter kilo of soil and three quarters of a kilo of ash**



**Figure 5: Non-plant growth when using quarter kilo of ash and three quarters of a kilo of soil**

**Table 1: Percentages of some elements in hospital waste ash**

Elements	Percentage
total nitrogen	0.189%
Total phosphorus	0.087%
total potassium	0.47%
organic matter	25.476%

In this study, the ash from burning hospital waste had a catastrophic effect on the fava bean plant. It stopped growing completely, and even when only small amounts were used, it still caused the plant to grow in a way that wasn't straight. This may be due to the fact that the ash from incinerating hospital waste contains very high levels of heavy metals that have a very toxic effect on plants and soil, as indicated by Honest *et al.*, [17], and Selman *et al.*, [18]. The ash from incinerating hospital waste contained high levels of heavy metals, including nickel, cobalt, and lead [19, 20]. Therefore, the results of the current study were clear that the ash from incinerating hospital waste is a toxic and lethal substance for plants, especially fava beans.

## CONCLUSIONS

It is concluded from the current study that contaminated and hazardous waste should be buried in relation to medical waste, and restrictions should be placed on access to the waste disposal site (building a fence around the place to prevent access by animals and children) and allocating closed places for all medical waste containers after sorting them instead of parking them in hospital facilities and corridors.

## REFERENCES

1. Kularatne, R. K. A. (2024). Biomedical waste generation at Ayurveda hospitals in South Asia: A mini review of the composition, quantities and characteristics. *Waste Management & Research*, 42(2), 95-110. <https://doi.org/10.1177/0734242X231178225>.
2. Ahmed, T. M. K., Yosif, K. M., & Mohammed, S. F. (2024). Identification and isolation of bacterial strains and study of their ability to degrade trimethoprim in the wastewater of some kirkuk hospitals. *S. Asian J. Life Sci*, 13, 8-12. <http://dx.doi.org/10.17582/journal.sajls/2024/13.8.12>.
3. Chandrappa, R., & Das, D. B. (2024). Biomedical Waste. In *Solid Waste Management: Principles and Practice* (pp. 243-297). Cham: Springer International Publishing.
4. Abu Qdais, H., Rabi, A. & Abdulla, F. (2007). Characteristics of the medical waste generated at the Jordanian hospitals. *Clean Techn Environ Policy* 9, 147–152. <https://doi.org/10.1007/s10098-006-0077-0>.
5. Ghanimeh, S., Gómez-Sanabria, A., Tsydenova, N., Štrbová, K., Iossifidou, M., & Kumar, A. (2019). Two-level comparison of waste management systems in low-, middle-, and high-income cities. *Environmental Engineering Science*, 36(10), 1281-1295. <https://doi.org/10.1089/ees.2019.0047>.
6. Padmanabhan, K. K., & Barik, D. (2019). Health hazards of medical waste and its disposal. In *Energy from toxic organic waste for heat and power generation* (pp. 99-118). Woodhead Publishing. <https://doi.org/10.1016/B978-0-08-102528-4.00008-0>.

7. Peng, J., Wu, X., Wang, R., Li, C., Zhang, Q., & Wei, D. (2020). Medical waste management practice during the 2019-2020 novel coronavirus pandemic: Experience in a general hospital. *American journal of infection control*, 48(8), 918-921. <https://doi.org/10.1016/j.ajic.2020.05.035>.
8. Vasistha, P., Ganguly, R., & Gupta, A. K. (2018). Biomedical waste generation and management in public sector hospital in Shimla City. In *Environmental Pollution: Select Proceedings of ICWEES-2016* (pp. 225-232). Springer Singapore. [https://doi.org/10.1007/978-981-10-5792-2\\_19](https://doi.org/10.1007/978-981-10-5792-2_19).
9. Jaber, S., Aljawad, A. A., Prisecaru, T., & Pop, E. (2021). The environmental situation of the ash medical waste in Baghdad city, Iraq. In *E3S Web of Conferences* (Vol. 286, p. 02017). EDP Sciences. <https://doi.org/10.1051/e3sconf/202128602017>.
10. Voshell, S., Mäkelä, M., & Dahl, O. (2018). A review of biomass ash properties towards treatment and recycling. *Renewable and Sustainable Energy Reviews*, 96, 479-486. <https://doi.org/10.1016/j.rser.2018.07.025>.
11. Wierzbowska, J., Sienkiewicz, S., Żarczyński, P., & Krzebietke, S. (2020). Environmental application of ash from incinerated biomass. *Agronomy*, 10(4), 482. <https://doi.org/10.3390/agronomy10040482>.
12. Petlickaitė, R., Jasinskas, A., Mioldažys, R., Romaneckas, K., Praspaliauskas, M., & Balandaitė, J. (2022). Investigation of pressed solid biofuel produced from Multi-Crop biomass. *Sustainability*, 14(2), 799. <https://doi.org/10.3390/su14020799>.
13. Petlickaitė, R., Jasinskas, A., Domeika, R., Pedišius, N., Lemanas, E., Praspaliauskas, M., & Kukharets, S. (2023). Evaluation of the Processing of Multi-Crop Plants into Pelletized Biofuel and Its Use for Energy Conversion. *Processes*, 11(2), 421. <https://doi.org/10.3390/pr11020421>.
14. Petlickaitė, R., Jasinskas, A., Venslauskas, K., Navickas, K., Praspaliauskas, M., & Lemanas, E. (2024). Evaluation of Multi-Crop Biofuel Pellet Properties and the Life Cycle Assessment. *Agriculture*, 14(7), 1162. <https://doi.org/10.3390/agriculture14071162>.
15. Ahmed, T. M. K., Mohammed, S. F., & Yosif, K. M. (2023). Biodegradation of Crude Oil Contaminated Soil in North Oil Refineries/Kirkuk/Iraq by Using *Acinetobacter* spp. and *Streptomyces* spp. *Journal of Survey in Fisheries Sciences*, 10(3S), 2286-2293.
16. Windfeld, E. S., & Brooks, M. S. L. (2015). Medical waste management—A review. *Journal of environmental management*, 163, 98-108. <https://doi.org/10.1016/j.jenvman.2015.08.013>.
17. Honest, A., Manyele, S. V., Saria, J. A., & Mbuna, J. (2020). Assessment of the heavy metal-levels in the incinerators bottom-ash from different hospitals in Dares Salaam. *African Journal of Environmental Science and Technology*, 14(11), 347-360. <https://doi.org/10.5897/AJEST2020.2891>.
18. Selman, H., Kubba, H., Al-Mukaram, N., & Alkateeb, R. (2021, March). Heavy metal pollution from hospital waste incinerators: A case study from Al-Muthanna province, Iraq. In *IOP Conference Series: Materials Science and Engineering* (Vol. 1090, No. 1, p. 012036). IOP Publishing. <https://doi.org/10.1088/1757-899X/1090/1/012036>.
19. Hussein, I. F., & Hussain, B. A. (2024, October). Health and Environmental Risk Assessment of Heavy Metals Contamination from Medical Wastes Ash of Baghdad Hospitals. In *Journal of Physics: Conference Series* (Vol. 2857, No. 1, p. 012049). IOP Publishing. <https://doi.org/10.1088/1742-6596/2857/1/012049>.
20. Amfo-Otu, R., Kyerewaa, S. G., Ofori, E. A., & Sadick, A. (2015). Comparative study of heavy metals in bottom ash from incinerators and open pit from healthcare facilities in Ghana. *Octa Journal of Environmental Research*, 3(1).