

Review Article

Role of Diet in Cancer Prevention

Dr. Anil Batta

Professor & Head, Department Of Medical Biochemistry Govt. Medical College, Amritsar, India

*Corresponding Author

Dr. Anil Batta

Article History: | Received: 05.03.2020 | Accepted: 26.04.2020 | Published: 30.04.2020 |

Abstract: Cancer chemoprevention is defined as the application of natural or synthetic agents to suppress or reverse cancer development and progression. In this field especially diet derived compounds have recently attracted researchers' attention as potential therapeutics generally exerting low toxicity compared with regular drugs. This review presents a survey of recent findings concerning the most promising dietary chemopreventive agents such as green tea polyphenols (i.e. catechins), long-chain polyunsaturated fatty acids, carotenoids, glucosinolates/isothiocyanates, vitamins (i.e. vitamin D and folate) and minerals (i.e. calcium and selenium). Molecular targets involved in intrinsic pathways affected by these natural compounds are also shortly discussed.

Keywords: cancer chemoprevention, green tea polyphenols, polyunsaturated fatty acids, glucosinolates, carotenoids.

Copyright © 2020: This is an open-access article distributed under the terms of the Creative Commons Attribution license which permits unrestricted use, distribution, and reproduction in any medium for non commercial use (NonCommercial, or CC-BY-NC) provided the original author and source are credited.

INTRODUCTION

As there is overall agreement that preventing disease is more desirable than curing, a multitude of potential chemopreventive compounds are under investigation. The term chemoprevention, first introduced by Sporn, means inhibiting, suppressing or reversing cancer development and progression with the use of natural or synthetic agents (Sporn, M. B. *et al.*, 1976, May). Chemopreventives should be characterized by low toxicity comparable with therapeutic drugs and a possibility to be taken orally. Nowadays, a new definition of chemoprevention can be stated and it can be described as the use of compounds suppressing or reversing the molecular pathways leading to cancer development and metastasis. Different dietary patterns in distinct countries are related to higher or lower incidences of cancer development in these ethnic groups, which confirms an inverse association between food borne preventive compounds and carcinogenesis. Lower incidence of breast, prostate and gastrointestinal tract cancer were always observed for Asian populations (Kurahashi, N. *et al.*, 2007; & Rose, D. P. 1997; & Dorai, T., & Aggarwal, B. B. 2004). Migration of people and adopting nutritional habits leads to alterations in cancer types and rates [WHO]. Diet derived compounds broadly studied for their chemopreventive activity include a wide group of polyphenols (i.e. green tea polyphenols, soy flavonoid genistein, quercetin from onions, resveratrol from grapes or curcumin), polyunsaturated fatty acids (n-3

PUFAs), carotenoids (i.e. β -carotene, lycopene or lutein), vitamins (D, E, C or folic acid) and minerals (Se, Zn, Ca) as well a dietary fiber. For many years it has been believed that diet can influence various diseases and the impact is often positive for the human organism. It was suggested already two and half thousand years ago by Hippocrates, who said "Let food be thy medicine and medicine be thy food". Diet can deliver both pro-healthy and harmful substances. Some can act as carcinogens and initiate or stimulate tumor growth. However, food includes an abundance of bioactive compounds which are beneficial to the human organism. There are data indicating a negative relationship between sufficient vegetable and fruit intake and cancer incidence. What is more, laboratory studies confirm this inverse association. It has been found that the risk of cancer incidence is half for people who eat at least five servings of fruits and vegetables per day (Surh, Y. J. 2003). A well-balanced diet should provide a rich source of macronutrients (omega-3 fatty acids, fiber) and micronutrients (i.e. selenium, calcium, vitamin D and E, folates) (Vaisman, N., & Arber, N. 2002; & Johnson, I. T. 2004), as well as phytochemicals (i.e. polyphenols, flavonoids, carotenoids) (Mehta, R. G. *et al.*, 2010), which exert cancer preventive activities. Various ways of administration are proposed to deliver such food derived compounds in populations with normal risk of cancer. They include simple daily consumption of fruits and vegetables or functional food enriched with several phytochemicals. Food extracts

can also be applied as food supplements (Mehta, R. G. *et al.*, 2010; & Kelloff, G. J. *et al.*, 2000).

Green Tea Polyphenols

It was estimated that mean consumption of tea per day is around 120 ml per person (McKay, D. L., & Blumberg, J. B. 2002). Green tea, always popular in China, Japan and India, now has gained popularity in many other countries. Black, oolong and green tea is derived from the same plant, *Camellia sinensis*, which is exposed to different processing methods before consumption. To obtain black tea the leaves of *Camellia sinensis* are fermented for a long period of time, which leads to formation of polymeric compounds, i.e. theaflavins and thearubigins. Oolong tea, often called half fermented, due to shorter fermentation time, contains polymeric as well as monomeric polyphenols. Non-fermented green tea is made from fresh tea leaves exposed to steaming and drying (to eliminate the oxidation process by inactivating the polyphenol oxidase) and contains an abundance of polyphenols naturally occurring in *Camellia sinensis* (Kelloff, G. J. *et al.*, 2000; & McKay, D. L., & Blumberg, J. B. 2002). Green tea contains proteins (including enzymes), amino acids, carbohydrates, lipids, vitamins (B, C, E) and minerals (i.e. Ca, Mg, Cr, Fe, Zn, F, K) (McKay, D. L., & Blumberg, J. B. 2002). Green tea has been found to exert a beneficial effect on the human organism and, although results are mixed, a number of studies have revealed a correlation with a reduction of chronic disease risk, including cancer (e.g. breast, esophageal, lung, colorectal, stomach, bladder, kidney, prostate, skin, pancreatic, ovarian) (McKay, D. L., & Blumberg, J. B. 2002), cardiovascular heart disease (coronary heart disease, hypertension, atherosclerosis) (Johnson, I. T. 2004), Parkinson disease (Kelloff, G. J. *et al.*, 2000), and Alzheimer disease (Kucuk, O. 2002). Green tea can also promote oral health (Kucuk, O. 2002; & Greenwald, P. *et al.*, 2002) and bone health (Kucuk, O. 2002; & Mehta, R. G. *et al.*, 2010). Consumption of this beverage is also being linked with anti-ageing processes, mainly due to its antioxidant properties (Kelloff, G. J. *et al.*, 2000; & McKay, D. L., & Blumberg, J. B. 2002). The positive health effect of green tea is mostly associated with the abundance of polyphenols, particularly flavonoids.

AS ANTIOXIDANT

Green tea polyphenols are responsible for significant and widely studied antioxidant properties of this beverage. Together with reactive oxygen and nitrogen species' scavenging ability, they exert beneficial effects on the human organism, protecting against harmful cigarette smoke, pollution or stress as well as cancer development (Kucuk, O. 2002; & Kurahashi, N. *et al.*, 2007). It has been revealed that catechins obtained from *Camellia sinensis* show higher antioxidant potential than generally known antioxidants, vitamin E and C (Mehta, R. G. *et al.*, 2010; & Dorai, T.,

& Aggarwal, B. B. 2004). Epigallocatechin gallate (EGCG) after epicatechin gallate (ECG) is the most effective of tea catechins. Their phenolic hydroxy groups on B and D rings are responsible for radical scavenging properties (McKay, D. L., & Blumberg, J. B. 2002). Moreover, they have the ability to chelate metal ions such as iron and copper, thus leading to decreased hydroxylic radical creation through inhibition of Fenton and Haber-Weiss reactions (Greenwald, P. *et al.*, 2002). Besides direct antiradical influence, they can also suppress redox-sensitive transcription factors or activate antioxidant enzymes, i.e. superoxide dismutases (Chen, C., & Kong, A. N. T. 2005).

Anticancer

Green tea polyphenols were shown to reduce the risk of a multitude of cancer types (see above). It is still not clear if the influence of green tea catechins on many molecular targets is attributed to their antioxidant activity or they directly and independently affect these cell signaling pathways (Rose, D. P. 1997). Polyphenols from *Camellia sinensis* may act through stimulation of apoptosis in cancer cells, both by inactivating anti-apoptotic (Bcl-2, Bcl-X1) and up-regulating pro-apoptotic (Bax, Bak, Bim, PUMA) proteins. Induced caspase-3 and -9 activities have also been shown (Kurahashi, N. *et al.*, 2007; Rose, D. P. 1997; & Dorai, T., & Aggarwal, B. B. 2004). The induction of cell cycle arrest in cancer cells by EGCG and other catechins has been broadly studied. It was reported that they stimulate the expression of p21 and p27 proteins, with simultaneous inhibition of cyclin D1, cyclin-dependent kinase 2 and cyclin-dependent kinase 4, thus leading to G1 growth arrest (Sporn, M. B. *et al.*, 1976; May, Kucuk, O. 2002; & Greenwald, P. *et al.*, 2002). Multiple signaling cascades engaged in cell survival, proliferation, differentiation and metabolism were found to be influenced by green tea polyphenol activity.

Antiangiogenic Activity

Creation of new blood vessels from pre-existing ones inside a tumor, called angiogenesis, promotes cancer development through its nourishment and oxygenation. Besides inhibiting activity of metalloproteinases which are involved in angiogenesis, EGCG was revealed to affect vascular endothelial growth factor (VEGF) binding ability to its receptor. VEGF is closely related to tumor-induced angiogenesis, acting as a mitogen factor for endothelial cells (Chen, C., & Kong, A. N. T. 2005; Kurahashi, N. *et al.*, 2007; & Rose, D. P. 1997).

Polyunsaturated Fatty Acids

A multitude of epidemiological studies have indicated an association between polyunsaturated fatty acid consumption and cancer risk, especially breast, prostate and colon cancer (Greenwald, P. *et al.*, 2002; & Mehta, R. G. *et al.*, 2010). A protective effect of the long-chain PUFAs is also revealed by *in vitro* and animal studies (Dorai, T., & Aggarwal, B. B. 2004).

The ratio of consumed n-6 to n-3 polyunsaturated fatty acids seems to play the major role in cancer development (Mehta, R. G. *et al.*, 2010). They are not synthesized in humans (essential fatty acids), so they must be supplemented with diet. The metabolic precursors of n-3 PUFAs [i.e. eicosapentaenoic acid (EPA), docosahexaenoic acid (DHA) and n-6 PUFAs

(i.e. arachidonic acid (AA), γ -linolenic acid (GLA)], are respectively α -linolenic acid (ALA, n-3) and linoleic acid (LA, n-6) (Mehta, R. G. *et al.*, 2010). Linoleic acid is found in large amounts in sunflower, soy and corn oils, while ALA is present in linseed and rapeseed oils. Long-chain fatty acids n-3 can be found in cold-water fish.



Fig.1 Food Pyramid In Cancer

Chemoprotective Activity Of N-3

Dietary PUFAs exert a suppressing (n-3) or promoting (n-6) effect on cancer mainly through their derivatives, eicosanoids, hormone-like lipids. Prostaglandins (PG), thromboxanes (TX) and leukotrienes (LT) are synthesized in enzymatic reactions with use of cyclooxygenases (COX) and lipoxygenases (LOX). The n-3 and n-6 long chain fatty acids compete for the enzymes and eicosanoids derived from them reveal opposing effects on cell proliferation, differentiation, angiogenesis, metastasis and inflammatory responses (Mehta, R. G. *et al.*, 2010; & Chen, C., & Kong, A. N. T. 2005). The first step of eicosanoid production is the hydrolytic release of PUFA from the cell membrane. Since arachidonic acid (AA, n-6) is the most abundant PUFA among phospholipids in

cell membranes, pro-inflammatory and cancer promoting eicosanoids are mainly produced. Diet enrichment with n-3 PUFAs may silence the effect of AA-derived eicosanoids through replacement of AA in membranes, thus increasing the amount of precursors for biosynthesis of cancer suppressing prostaglandins and leukotrienes (Chen, C., & Kong, A. N. T. 2005; & Rose, D. P. 1997). Moreover, high consumption of n-3 long chain fatty acids leads to decreased production of AA from LA, due to stronger affinity of n-3 than n-6 PUFAs to enzymes involved in this process (Mehta, R. G. *et al.*, 2010; & Rose, D. P. 1997). It was revealed that PUFAs and their derivatives may influence several molecular signaling pathways involved in basic processes including cell growth, differentiation, apoptosis or metastasis.




Fig-2.Colorectal Cancer

Glucosinolates/Isothiocyanates

Crucifers (the plant family *Cruciferae*, also known as *Brassicaceae*) include several widely

consumed vegetables, e.g. cabbage, broccoli, cauliflower, parsnip, Brussels sprouts, radish, horseradish, turnip and kale Chen, C., & Kong, A. N. T.

2005; & Kurahashi, N. *et al.*, 2007). Epidemiological data suggest an inverse association between cruciferous vegetables intake and risk of many cancers, e.g. lung, breast, stomach, pancreas, prostate, bladder, thyroid, skin and colon cancer. Significantly, this correlation was found to be more prominent than the general effect linked to vegetable and fruit consumption (Kurahashi, N. *et al.*, 2007; & Rose, D. P. 1997). Despite the variety of bioactive compounds present in crucifers, i.e. polyphenols, minerals and vitamins, a superior role in cancer prevention is ascribed to glucosinolates and their decomposition products, isothiocyanates (Chen, C., & Kong, A. N. T. 2005; & Dorai, T., & Aggarwal, B. B. 2004). Glucosinolates (GS) characterized to date are a well-defined group of chemical compounds consisting of three basic structures including a β -D-thioglucose group, sulfonated oxime group and side chain derived from one of the protein amino acids (Dorai, T., & Aggarwal, B. B. 2004; & Surh, Y. J. 2003). Due to the acidic nature of sulfate groups, in plants glucosinolates are accumulated as potassium salts sequestered to vacuoles and probably protect against herbivores and microbes (Kurahashi, N. *et al.*, 2007). Due to the acidic nature of sulfate groups, in plants glucosinolates are accumulated as potassium salts sequestered to vacuoles and probably protect against herbivores and microbes (Chen, C., & Kong, A. N. T. 2005). Glucosinolates are highly stable and when consumed cannot act as bioactive compounds until subjected to enzymatic hydrolysis due to myrosinase action. Several mechanisms of cancer protection by glucosinolates/isothiocyanates have been proposed. The most studied is the influence of the compounds on detoxification pathways leading to increased activity of phase I and phase II enzymes and resulting in lowered amounts of carcinogens within the body due to their decreased formation as well as increased excretion.

Vitamins And Minerals

Intake of minerals and vitamins in daily diet is essential for health maintenance. Although they are present in trace amounts within tissues, they play a significant role in a variety of biological processes. Some of them also have a positive impact on the process of carcinogenesis. Anticancer activity of vitamins E and C was observed, mostly due to their antioxidant effects.

Vitamin D

Vitamin D is a fat-soluble prohormone naturally occurring in very small amounts in food. Two forms of this vitamin can be distinguished: calciferol (D_2) and cholecalciferol (D_3), which is produced in skin after exposure to UV-B light. Active forms of vitamin D are obtained in the body after their conversion in the liver to calcidiol (25-hydroxyvitamin D) and in kidneys to calcitriol (1, 25-dihydroxyvitamin D) (Rose, D. P. 1997). It has been revealed that human cells, both normal and cancer cells, demonstrate specific receptors (VDR) for vitamin D metabolites, thus being

susceptible to their activity (Rose, D. P. 1997; & Dorai, T., & Aggarwal, B. B. 2004). Vitamin D action has been attributed to the regulation of multiple genes (more than 60) engaged in cell proliferation, differentiation, apoptosis and angiogenesis (Rose, D. P. 1997; & Dorai, T., & Aggarwal, B. B. 2004). Biological activity of calcitriol is exerted when it binds to VDR receptors, which leads to formation of a heterodimer with retinoid X receptor and expression of specific genes containing vitamin D response element.

Folate

Folic acid (also known as vitamin B_9) is a constituent found in large amounts in green leafy vegetables and fresh fruits. It plays a crucial role in proper cell division and growth and takes part in nucleotide biosynthesis [10]. Its cancer preventive effects are mainly attributed to the delivery of methyl groups for DNA synthesis and gene expression, thus protecting against DNA damage. It was suggested that insufficient folic acid consumption may trigger loss of DNA integrity through misincorporation of uracil to the chain of DNA and induces breaks in chromosomes. Reversion of these defects was shown after folate supplementation (Dorai, T., & Aggarwal, B. B. 2004; & Surh, Y. J. 2003). A reverse association was observed between folic acid intake and colon and breast cancer incidence. Animals fed with a diet poor in folate were more likely to suffer from colorectal cancer.

Selenium

A variety of *in vitro*, animal and epidemiological studies suggest a preventive role of selenium against human cancer development, especially tumors of the gastrointestinal tract. Low levels of selenium in serum were associated with increased risk of prostate cancer. It is suggested that selenium may inhibit initiation and promotion phases of carcinogenesis (Vaisman, N., & Arber, N. 2002). Besides its antioxidant effect on lipid peroxidation, it can alter the process of DNA adduct formation. It blocks formation of carcinogens and their binding to DNA. Selenium may also reduce cancer progression through improving the immune response in the organism, e.g. by stimulation of cytotoxic activity of natural killer cells (Surh, Y. J. 2003). Several studies have revealed selenium-induced apoptosis in tumor cells (Surh, Y. J. 2003; & Vaisman, N., & Arber, N. 2002) as well as inhibition of cell proliferation involving alternations in MAPK kinases and Akt kinase signaling pathways (Mehta, R. G. *et al.*, 2010; & Rose, D. P. 1997). It was also proposed that it could act as a chemopreventive agent by decreasing angiogenesis within the tumor (Chen, C., & Kong, A. N. T. 2005; & Dorai, T., & Aggarwal, B. B. 2004). What is interesting, small amounts of Se added to a diet deficient in this micronutrient results in improved activity of selenoproteins and detoxification enzymes of phase I and II as well as ameliorated function of the immune system, whereas supranutritional intake of

selenium leads to cancer prevention through the above-described influence on the cell cycle, apoptosis and angiogenesis (Chen, C., & Kong, A. N. T. 2005; & Rose, D. P. 1997).

Calcium

Calcium constitutes an essential macroelement for human health. Dairy products (milk, yogurts, and cheese), dark green vegetables, legumes and nuts constitute a good source of this micronutrient. Although some results are inconsistent, a number of epidemiological studies have attributed calcium intake with decreased colon adenoma and cancer (Dorai, T., & Aggarwal, B. B. 2004; & Surh, Y. J. 2003). The mechanism underlying this protective effect is mainly due to binding and inactivating the pro-carcinogenic activity of bile acids on colon epithelial cells. In addition, calcium was found to inhibit epithelial cell proliferation or induce their differentiation through altering cell signaling pathways (Kurahashi, N. *et al.*, 2006; Rose, D. P. 1997; Dorai, T., & Aggarwal, B. B. 2004; & Surh, Y. J. 2003).

CONCLUSION

Based on various studies outlined in this review reporting a broad spectrum of naturally occurring compounds with chemopreventive activity, intense consideration of them as potential therapeutics is quite intelligible. Some of these may be already found as extracts or pure substances available for daily supplementation and are becoming a common dietary habit. However, providing a well-balanced diet containing an abundance of biologically active compounds should become a daily habit.

REFERENCES

1. Chen, C., & Kong, A. N. T. (2005). Dietary cancer-chemopreventive compounds: from signaling and gene expression to pharmacological effects. *Trends in pharmacological sciences*, 26(6), 318-326.
2. Dorai, T., & Aggarwal, B. B. (2004). Role of chemopreventive agents in cancer therapy. *Cancer letters*, 215(2), 129-140.
3. Greenwald, P., Milner, J. A., Anderson, D. E., & McDonald, S. S. (2002). Micronutrients in cancer chemoprevention. *Cancer and metastasis Reviews*, 21(3-4), 217-230.
4. Johnson, I. T. (2004). New approaches to the role of diet in the prevention of cancers of the alimentary tract. *Mutation Research/Fundamental and Molecular Mechanisms of Mutagenesis*, 551(1-2), 9-28.
5. Kelloff, G. J., Crowell, J. A., Steele, V. E., Lubet, R. A., Malone, W. A., Boone, C. W., ... & Ali, I. (2000). Progress in cancer chemoprevention: development of diet-derived chemopreventive agents. *The Journal of nutrition*, 130(2), 467S-471S.
6. Kucuk, O. (2002). Cancer chemoprevention. *Cancer and Metastasis Reviews*, 21(3-4), 189-197.
7. Kurahashi, N., Iwasaki, M., Sasazuki, S., Otani, T., Inoue, M., & Tsugane, S. (2007). Soy product and isoflavone consumption in relation to prostate cancer in Japanese men. *Cancer Epidemiology and Prevention Biomarkers*, 16(3), 538-545.
8. McKay, D. L., & Blumberg, J. B. (2002). The role of tea in human health: an update. *Journal of the American College of Nutrition*, 21(1), 1-13.
9. Mehta, R. G., Murillo, G., Naithani, R., & Peng, X. (2010). Cancer chemoprevention by natural products: how far have we come?. *Pharmaceutical research*, 27(6), 950-961.
10. Rose, D. P. (1997). Dietary fatty acids and cancer. *The American journal of clinical nutrition*, 66(4), 998S-1003S.
11. Sporn, M. B., Dunlop, N. M., Newton, D. L., & Smith, J. M. (1976, May). Prevention of chemical carcinogenesis by vitamin A and its synthetic analogs (retinoids). In *Federation proceedings* (Vol. 35, No. 6, pp. 1332-1338).
12. Surh, Y. J. (2003). Cancer chemoprevention with dietary phytochemicals. *Nature Reviews Cancer*, 3(10), 768-780.
13. Vaisman, N., & Arber, N. (2002). The role of nutrition and chemoprevention in colorectal cancer: from observations to expectations. *Best Practice & Research Clinical Gastroenterology*, 16(2), 201-217.