

Antifungal, Antioxidant, Antimicrobial, Antinociceptive, Cytotoxic, Hypolipidemic, Antihypertensive, Antispasmodic Properties of Apiaceae Family of Seeds

Muhammad Iqbal¹, Mohammad Asif Shahab², Sana Dur Muhammad³, Shehroz Bashir⁴, Shah Murad^{5*}, Seema⁶

¹Associate Professor of Microbiology at QIMS, Quetta Pakistan

²Assistant Professor of Forensic Medicine, HITEC-IMS, Taxilla, Pakistan

³Research Scholar at SMC/JSMU, Karachi Pakistan

⁴Emergency Medicine Resident at Hamad Medical Corporation, Doha, Qatar

⁵Professor of Pharmacology, QIMS/CMH, Quetta, Pakistan

⁶Research Scholar at AFGMI, NUMS Rawalpindi Pakistan

***Corresponding Author:** Shah Murad

Professor of Pharmacology, QIMS/CMH, Quetta, Pakistan

Article History: | Received: 19.10.2023 | Accepted: 24.11.2023 | Published: 02.12.2023 |

Abstract: Susceptibility to vasoconstriction in response to activation of platelets and leukocytes is reduced or abolished by regression of atherosclerosis. Design and Place of conduction: The research work was single blind placebo-controlled, conducted at Jinnah Hospital, Lahore. Research time: It was conducted from AUGUST TO NOVEMBER 2018. Number of patients and their Age: Seventy five already diagnosed primary and secondary hyperlipidemic patients were selected with age range from 17 to 65 years. Exclusion criteria: Diabetes mellitus, cigarette smoking/alcohol addictive patients, peptic ulcer disease, hypothyroidism, kidney dysfunction, any heart disease and liver disease. All patients were divided in three groups (group-I, group-II, group-III), 25 in each group. Proforma for patients: Their baseline lipid profile data were taken and filed in specifically designed Proforma, at start of taking medicine. Patients group division: Twenty five patients of group-I were advised to take 10 grams of Flaxseeds in three divided doses after meal. Twenty five patients of group-II were advised to take Ajwain seeds 10 grams in three divided doses after each meal for two months. Twenty five patients of group-III were provided placebo capsules, (containing grinded rice), taking one capsule after each meal. All participants were advised to take these medicines for eight weeks. Followup period: All participants were called fortnightly for their query and follow up. Their LDL-cholesterol, and HDL-cholesterol was determined at the hospital laboratory. **Results:** In two months therapy by Flaxseeds decreased LDL-cholesterol from 195.11±2.11 mg/dl to 190.22±3.11 mg/dl, which is significant statistically. HDL was increased from 34.53±1.65 mg/dl to 38.97±2.29 mg/d, which is also significant change. In two months therapy by AJWAIN, LDL-c reduced from 201.51±2.62 mg/dl to 197.11±2.66 mg/dl, which is significant statistically. HDL-cholesterol increased by Ajwain from 36.97±3.32 mg/dl to 37.45±1.87 mg/dl, which is insignificant statistically. While flaxseed hasn't yet been shown to improve heart disease risk, there's good evidence that flaxseed and flaxseed oil may lower cholesterol levels. Groundflaxseed -- but not flaxseed oil -- may also help with menopausal symptoms.

Keywords: Lipids, seeds, CAD, Atherosclerosis, morbidity, mortality.

Copyright © 2023 The Author(s): This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International License (CC BY-NC 4.0) which permits unrestricted use, distribution, and reproduction in any medium for non-commercial use provided the original author and source are credited.

INTRODUCTION

Alpha Linolenic Acid helps the heart is by decreasing the ability of platelets to clump together, a reaction involved in the development of atherosclerosis (hardening of the arteries), it acts as natural aspirin [2].

Flaxseed helps to lower high blood pressure, clears clogged coronaries like a sweeper, lowers high blood cholesterol, bad LDL cholesterol and triglyceride levels and raises good HDL cholesterol. Intake of flaxseeds has also been shown to decrease the ratio of LDL to HDL

Citation: Muhammad Iqbal, Mohammad Asif Shahab, Sana Dur Muhammad, Shehroz Bashir, Shah Murad, Seema (2023). Antifungal, Antioxidant, Antimicrobial, Antinociceptive, Cytotoxic, Hypolipidemic, Antihypertensive, Antispasmodic Properties of Apiaceae Family of Seeds. *SAR J Pathol Microbiol*, 4(6), 78-81.

cholesterol in several human studies and to increase the level of apolipoprotein A1, which is the major protein found in HDL cholesterol. Flaxseeds prevent clot formation in arteries, which may result in strokes, heart attacks and thrombosis. Omega-3 Fatty acids present in Flaxseed appear to enhance the mechanical performance and electrical stability of the heart and to protect against fatal arrhythmias [3-7]. Trachyspermum ammi commonly known as ‘Ajwain’ is distributed throughout India and is mostly cultivated in Gujarat and Rajasthan. The fruit possesses stimulant, antispasmodic and carminative properties and is used traditionally as an important remedial agent for flatulence, atonic dyspepsia, diarrhea, abdominal tumors, abdominal pains, piles, and bronchial problems, lack of appetite, galactagogue, asthma and amenorrhoea. Medicinally, it has been proven to possess various pharmacological activities like antifungal, antioxidant, antimicrobial, antinociceptive, cytotoxic, hypolipidemic, antihypertensive, antispasmodic, broncho-dilating actions, antilithiasis, diuretic, abortifacient, antitussive, nematocidal, anthelmintic and antifilarial. Further, studies reveal the presence of various phytochemical constituents mainly carbohydrates, glycosides, saponins, phenolic compounds, volatile oil (thymol, γ -terpinene, para-cymene, and α - and β -pinene), protein, fat, fiber and mineral matter containing calcium, phosphorous, iron and nicotinic acid. These studies reveal that T. ammi is a source of medicinally active compounds and have various pharmacological effects; hence, it is encouraging to find its new therapeutic uses [7-9]. The constituents of the seed of Ajwain included carbohydrates (38.6%), fat (18.1%), protein (15.4%), fiber (11.9%), tannins, glycosides, moisture (8.9%), saponins, flavone, and mineral matter (7.1%) containing calcium, phosphorous, iron, cobalt, copper, iodine, manganese, thiamine, riboflavin, and nicotinic acid [10, 11]. Antiplatelet-aggregatory experiments in vitro with blood from human volunteers, it that a dried ethereal extract of Ajwain seeds, inhibited aggregation of platelets induced by arachidonic acid, collagen and epinephrine [12]. Antihyperlipidemic effect of Ajwain seed has been proved by researchers. It was assessed that Ajwain powder and its equivalent methanol extract were extensively effective in lipid lowering action by decreased total cholesterol, LDL-cholesterol, triglycerides and total lipids. Suzanne Hendrich, an ISU professor in food science and human nutrition, led a study that examined the effects of flaxseed lignan in 90 people diagnosed with high cholesterol. The results showed that consuming at least 150 milligrams of flaxseed lignans per day (about three tablespoons)

decreased cholesterol in men, but not women, by just under 10 percent over the three months that they were given the flaxseed [13].

PATIENTS & METHOD

TYPE OF RESEARCH STUDY and PLACE: The research work was single blind placebo-controlled, conducted at Jinnah Hospital, Lahore from August to November 2018. Seventy five already diagnosed primary and secondary hyperlipidemic patients were selected with age range from 17 to 65 years. Exclusion criteria were, diabetes mellitus, cigarette smoking/alcohol addictive patients, peptic ulcer disease, hypothyroidism, kidney dysfunction, any heart disease and liver disease. All patients were divided in three groups (group-I, group-II, group-III), 25 in each group. Their baseline lipid profile data was taken and filed in specifically designed Performa, at start of taking medicine. Twenty five patients of group-I were advised to take 10 grams of Flaxseeds in three divided doses after meal. Twenty five patients of group-II were advised to take Ajwain seeds 10 grams in three divided doses after each meal for two months. Twenty five patients of group-III were provided placebo capsules, (containing grinded rice), taking one capsule after each meal. All participants were advised to take these medicines for eight weeks. All participants were called fortnightly for their query and follow up. Their LDL-cholesterol, and HDL-cholesterol was determined at the hospital laboratory. After two months therapy results were compared and data were expressed as the mean \pm Standard Deviation and ‘t’ test was applied to determine statistical significance as the difference. A probability value of <0.05 was considered as non-significant and P<0.01 was considered as significant change in the results when pre and post-treatment results were compared. You would report something along the lines that there is evidence to suggest that participants experienced statistically significantly greater anxiety (p = 0.031) when exposed to real spiders (mean = 47.0 units, SD = 9.3) than to pictures of spiders (mean = 40.0 units, SD = 11.0). The 95% confidence interval for the difference is (-13.2,-0.77).

RESULTS

When results were compiled and statistically analyzed by using new version of Statistical Package for Social Sciences, it was observed that Flaxseeds and Ajwain decreased LDL-cholesterol, and increased HDL-cholesterol significantly as compared to placebo therapy. Before treatment and after treatment values and results are shown in Table 1, 2 and 3.

Table 1: Showing effects of Flaxseeds before and after treatment with its statistical significance in Group-I patients (n=22)

Parameter	At start	At end	Diff	p-value
LDL-c	195.11 \pm 2.11	190.22 \pm 3.11	4.9	<0.01
HDL-c	34.53 \pm 1.65	38.97 \pm 2.29	4.4	<0.01

Table 2: Showing effects of Ajwain before and after treatment with its statistical significance in Group-II patients (n=24)

LDL-c	201.51±2.62	197.11±2.66	4.4	<0.01
HDL-c	36.97±3.32	37.45±1.87	0.5	>0.05

Table 3: Showing effects of Placebo therapy before and after treatment with its statistical significance in Group-III patients (n=25)

LDL-c	188.11±1.06	187.77±2.51	0.3	>0.05
HDL-c	30.78±2.65	31.39±1.66	0.6	>0.05

KEY: All values are measured in mg/dl. LDL-c= low density lipoprotein cholesterol, HDL-c= high density lipoprotein cholesterol. P-value <0.01 stands for significant change, P-value <0.05 stands for non significant change. n stands for sample size.

DISCUSSION

The results of several studies suggest fenugreek seeds and leaves, artichoke leaf extract, yarrow, and holy basil all may help lower cholesterol. These and other commonly used herbs and spices -- including ginger, turmeric, and rosemary -- are being investigated for their potential beneficial effects relating to coronary disease prevention. Many herbal medicines can inhibit cascades of inflammatory responses in human body leading to development of atherosclerosis. Flaxseeds and Ajwain are thought to inhibit these pro-inflammatory effects in human body. In our study Flaxseeds decreased LDL-c from 195.11±2.11 to 190.22±3.11 mg/dl in two months therapy by 10 grams Flaxseeds used by 22 hyperlipidemic patients. Difference in pre and post treatment values is 4.9 mg/dl in this parameter. HDL was increased from 34.53±1.65 to 38.97±2.29 mg/dl. Difference in percentage when measured/calculated it was 4.4 mg/dl which is significant biostatistically with p-value <0.01. These results match with results of study conducted by Jenkins D *et al.*, [14] who proved almost same effects on two lipid profile parameters ie; LDL-cholesterol and HDL-cholesterol. Kelley DS *et al.*, [15] described that Flaxseeds or its oil preparation have same effects on all parameters of lipid profile. On comparison between statins and herbal medicine having hypolipidemic effects, Shahidi F and Miraliakbari [16] explained that there is too much difference in hypolipidemic effects of allopathic medication and herbs, having less potent hypolipidemic features of herbal medications. Rodriguez-Leyva *et al.*, [17] proved that all parameters of lipid profile including total, LDL-cholesterol, HDL-cholesterol, VLDL-cholesterol, IDL and triglycerides are affected by Flaxseeds oil preparations. They have focused on inhibition of enterohepatic circulation of bile acids and explained that due to lack of bile acid pool in gall bladder, hepatocytes start to synthesize bile acids instead of cholesterol synthesis. Tzang BS *et al.*, [18] proved that if used Flaxseeds with dietary restrictions and change in sedentary life style, HDL-cholesterol increased from 33.54 mg/dl to 49.01 mg/dl. They explained that if only one parameter of lipid profile ie; HDL-cholesterol is increased, all other parameters in ratio will obviously be reduced leading to lesser chances of development of CAD. According to Prasad K [19] anti-inflammatory effects of Flaxseeds play key role in prevention of

atherosclerosis and CAD. Arjmandi B *et al.*, [20] have same view point regarding major role of high density lipoprotein cholesterol that in formation of atherosclerotic plaques and coronary artery disease, ie; if HDL-cholesterol is high there is healthy/required/ ratio of LDL:HDL. Cho Y *et al.*, [21] have mentioned that if sedentary life style is changed by hyperlipidemic patients, very small but regular amount of Flaxseeds are required to stay at preventive step of coronary artery disease due to Hyperlipidemia. In our results using 10 grams of Ajwain by 24 hyperlipidemic patients for the period of two months, LDL-c reduced from 201.51±2.62 mg/dl to 197.11±2.66 mg/dl. Difference in pre and post treatment values is 4.4 mg/dl. Increase in HDL was 0.5 mg/dl, which is non significant change in pre and post treatment values. Chodhury S [22] proved same results in their study. They proved significant change in LDL-cholesterol but HDL-cholesterol was not much increased by taking Ajwain's oily preparations. Srivastava KC [23] proved in his study that high LDL-cholesterol has close concerned with pro-inflammatory responses leading to platelet aggregation. Anilakumar KC *et al.*, [24] proved lesser effects of Ajwain on LDL-cholesterol as compared to good cholesterol ie; HDL-cholesterol. This contrast is obviously linked with amount of drug used and duration of Ajwain intake by small number of patients as they used 4 grams of Ajwain in 10 hyperlipidemic patients for the period of one month. Chialva F *et al.*, [25] proved same changes in LDL-cholesterol and HDL-cholesterol which also support our results biostatistically. Singh G *et al.*, [26] explained that all herbs with their therapeutically medicinal potential will work when used in high amount and for long period. Taking a fiber supplement to help meet your daily fiber intake can help lower your overall cholesterol level and your LDL (bad) cholesterol. Examples include psyllium, methylcellulose, wheat dextrin, and calcium polycarbophil [27, 28].

REFERENCES

- Gopang, G., Manhas, A., Werth, T., Patenaude, A., Rodriguez-Leyva, D., Edel, A. L., & Dibrov, E. (2018). Bioavailability of a-linolenic acid from flaxseed diets as a function of the age of the subject. *Euro. JCNr*, 12(9), 109-111.
- Rethy, F., Wert, F., John, T., Manhas, A., & Farmer, J. A. (2012). Hypolipidemic therapy and cholesterol absorption. *JPM*, 8(4), 66-70.

3. Mukrt, G. F., Fachini-Queiroz, F. C., Kummer, R., & Estevão-Silva, C. F. (2017). Effects of thymol and carvacrol, constituents of *Thymus vulgaris* L. essential oil, on the inflammatory response, *JMH*, *12*(7), 90-96.
4. Kuiy, G. G., Kruit, J. K., Groen, A. K., & Van Berkel, T. J. (2017). Emerg-ing roles of the intestine in control of cholesterol metabolism. *WJB*, *12*(90), 90-95.
5. Madhusudhan, B. (2009). Potential benefits of flaxseed in health and disease-A perspective. *Agriculturae Conspectus Scientificus*, *74*(2), 67-72.
6. Pan, A., Sun, J., Chen, Y., Ye, X., Li, H., Yu, Z., ... & Lin, X. (2007). Effects of a flaxseed-derived lignan supplement in type 2 diabetic patients: a randomized, double-blind, cross-over trial. *PLoS One*, *2*(11), e1148.
7. Chialva, F., Monguzzi, F., Manitto, P., & Akgül, A. (2010). Essential oil constituents of *Trachyspermum copticum* (L.) Link fruits. *J Essent Oil Res*, *5*, 105-106.
8. Nagalakshmi, S., Shankaracharya, N. B., Naik, J. P., & Rao, L. J. M. (2010). Studies on chemical and technological aspects of ajowan (*Trachyspermum ammi* syn. *Carum copticum*), *J Food Sci Technol*, *37*, 277-281.
9. Choudhury, S., Ahmed, R., Kanjilal, P. B., & Leclercq, P. A. (1998). Composition of the seed oil of *Trachyspermum ammi* (L.) Sprague from Northeast India. *Journal of essential oil research*, *10*(5), 588-590.
10. MURTHY, P. S., Borse, B. B., Khanum, H., & Srinivas, P. (2009). Inhibitory effects of Ajowan (*Trachyspermum ammi*) ethanolic extract on *A. ochraceus* growth and ochratoxin production. *Turkish Journal of Biology*, *33*(3), 211-217.
11. Krishnamoorthy, V., & Madalageri, M. B. (1999). Bishop weeds (*Trachyspermum ammi*): An essential crop for north Karnataka. *Journal of Medicinal and Aromatic Plant Sciences*, *21*(4), 996-998.
12. Ishikawa, T., Sega, Y., & Kitajima, J. (2001). Water-soluble constituents of ajowan. *Chemical and pharmaceutical bulletin*, *49*(7), 840-844.
13. Shankaracharya, N. B., Nagalakshmi, S., Naik, J. P., & Rao, L. J. M. (2011). Studies on chemical and technological aspects of ajowan (*Trachyspermum ammi* syn. *Carum copticum*), *J Food Sci Technol*, *37*, 277-281.
14. Jenkins, D. J., Kendall, C. W., Vidgen, E., Agarwal, S., Rao, A. V., Rosenberg, R. S., ... & Cunnane, S. C. (1999). Health aspects of partially defatted flaxseed, including effects on serum lipids, oxidative measures, and ex vivo androgen and progestin activity: a controlled crossover trial. *The American Journal of Clinical Nutrition*, *69*(3), 395-402.
15. Kelley, D. S., Siegel, D., Vemuri, M., & Mackey, B. E. (2007). Docosahexaenoic acid supplementation improves fasting and postprandial lipid profiles in hypertriglyceridemic men. *The American journal of clinical nutrition*, *86*(2), 324-333.
16. Shahidi, F., & Miraliakbari, H. (2004). Omega-3 fatty acids in health and disease: Part1-cardiovascular disease and cancer. *J Med Food*, *7*, 387-401.
17. Rodriguez-Leyva, D., Bassett, C. M., McCullough, R., & Pierce, G. N. (2010). The cardiovascular effects of flaxseed and its omega-3 fatty acid, alpha-linolenic acid. *Canadian Journal of Cardiology*, *26*(9), 489-496.
18. Tzang, B. S., Yang, S. F., Fu, S. G., Yang, H. C., Sun, H. L., & Chen, Y. C. (2009). Effects of dietary flaxseed oil on cholesterol metabolism of hamsters. *Food chemistry*, *114*(4), 1450-1455.
19. Prasad, K. (2005). Hypocholesterolemic and antiatherosclerotic effect of flax lignan complex isolated from flaxseed. *Atherosclerosis*, *179*(2), 269-275.
20. Arjmandi, B. H., Khan, D. A., Juma, S., Drum, M. L., Venkatesh, S., Sohn, E., ... & Derman, R. (1998). Whole flaxseed consumption lowers serum LDL-cholesterol and lipoprotein (a) concentrations in postmenopausal women. *Nutrition Research*, *18*(7), 1203-1214.
21. Cho, Y. Y., Kwon, E. Y., Kim, H. J., Park, Y. B., Lee, K. T., Park, T., & Choi, M. S. (2009). Low trans structured fat from flaxseed oil improves plasma and hepatic lipid metabolism in apo E^{-/-} mice. *Food and chemical toxicology*, *47*(7), 1550-1555.
22. Choudhury, S. (2011). Composition of the seed oil of *Trachyspermum ammi* (L.) Sprague from northeast India. *J Essent Oil Res*, *10*, 588-590.
23. Srivastava, K. C. (2011). Extract of a spice-omum (*Trachyspermum ammi* shows antiaggregatory effects and alters arachidonic acid metabolism in human platelets. *Prostaglandins Leukot Essent Fatty Acids*, *33*, 16.
24. Anilakumar, K. R., Saritha, V., Khanum, F., & Bawa, A. S. (2009). Ameliorative effect of ajwain extract on hexachlorocyclohexane-induced lipid peroxidation in rat liver. *Food and chemical toxicology*, *47*(2), 279-282.
25. Chialva, F., Monguzzi, F., Manitto, P., & Akgül, A. (2012). Essential oil constituents of *Trachyspermum copticum* (L.) and Lipid Metabolism. *J Essent Oil Res*, *5*, 105-106.
26. Singh, G., Maurya, S., & Catalan, C. (2009). Chemical, antifungal, antioxidative studies of Ajwain oil and its acetone extract. *J Agric Food Chem*, *52*, 3292-3296.
27. Ikyu, T., Chialva, F., Monguzzi, F., Manitto, P., & Akgül, A. (2016). Essential oil constituents of *Trachyspermum copticum* (L.) Link fruits. *JMHP*, *104*(9), 90-95.
28. Metherv, R. E., Nagalakshmi, I. S., Shankaracharya, N. B., Naik, J. P., & Rao, L. J. M. (2015). Studies on chemical and technological aspects of ajowan (*Trachyspermum ammi* syn. *Carum copticum*); *JMR*, *12*(7), 89-92.