

Review Article

Cosmetics' Safety: Gray Areas with Darker Inside-Mini Review

Abdul Kader Mohiuddin*

Assistant Professor, Department of Pharmacy, World University of Bangladesh

***Corresponding Author**

Abdul Kader Mohiuddin

Article History

Received: 03.08.2019

Accepted: 26.08.2019

Published: 30.09.2019

Abstract: Cosmetic items contain a wide scope of chemicals to which we are exposed every day. All cosmetics were separated into 3 classifications: rinse-off items (shower gel, shampoo, toothpaste, liquid soap, private soap, shaving foam) leave-on items (body cream, face cream, hand cream, antiperspirant, sunscreen, post-shaving astringent) and make-up ones (lipstick, lip-balm, foundation, nail polish). The utilization of a significant number of these substances is permitted inside specific limits, because of their toxicity at higher concentrations. Other significant viewpoints ought to be considered as, for example, the likelihood of long-term effects. In the light of the continuous and close nature of the contact on skin and mucosa with these items, it is significant that they don't contain conceivably perilous substances. Additionally, the everyday use and continuous exposition of people to a wide scope of personal consideration items and to various types of chemicals, got from a few sources, may cause the alleged "cocktail effect" because of the synergistic interaction of various substances and, likewise, the "added substance effect" in light of the nearness of a similar ingredient in numerous items. Actually, every one of the ingredients utilized in cosmetic items meet certain regulatory prerequisites. Notwithstanding, the utilization of numerous substances is permitted inside specific limits, because of their toxicity at higher concentrations.

Keywords: Safety issues of cosmetics; personal care products; sensitizations; skin irritation; contact dermatitis; formaldehyde releasers; phthalates family; parabens



Fig-1: Can Makeup Harm My Vision? [1,2].

Eye cosmetics are frequently applied among female populations of all age groups around the world. However, the migration of cosmetic products across the eyelid margin has been reported, and this is thought to exacerbate tear film instability and symptoms of dry eye. Makeup application and allergic reactions can cause significant eye injuries, and it's important to recognize what to avoid.

INTRODUCTION

Cosmetics and personal care products are ubiquitous. Currently, an increasing number of compounds are being assimilated in the formulation of cosmetic products as preservatives, fragrances, surfactants, etc. to intensify the performance, quality, value, and lifespan of cosmetics. Nevertheless, many of these chemical additives pose toxic effects to the human body, exhibiting health risks from a mild hypersensitivity to life-threatening anaphylaxis or lethal intoxication [3]. Most cosmetic products are directly applied on the skin and their ingredients can cross the cutaneous barrier to reach the systemic circulation [4]. With continued consumer concern and

Copyright @ 2019: This is an open-access article distributed under the terms of the Creative Commons Attribution license which permits unrestricted use, distribution, and reproduction in any medium for non commercial use (NonCommercial, or CC-BY-NC) provided the original author and source are credited.

several recent cosmetic-related public health controversies, the FDA should be given more resources and broader authority to protect consumer safety [5]. Toklu *et.al*, 2019 reported that more than 95% of cutaneous complaints are burning and itching. Interestingly, 60% of the cosmetic-related injuries did not consider any type of consultation [6]. The condition is found in more than 50% of women and 40% of men, creating a sizable demand for products designed to minimize skin sensitivity [7,8].

Moreover, such effects are underestimated because of the absence of formal and reliable monitoring systems ('cosmetovigilance') [9]. Fragranced ingredients are widespread diffused in cosmetic products but many of these may cause sensitizations, allergies and skin irritations [10]. A Senegalese cohort study of 147 women showed a statistically significant increase in the risk of hypertension and diabetes linked to the use of skin-lightening agents [11]. Both kojic acid and hydroquinone used as skin lighteners cause dryness of skin, peeling, and contact dermatitis [12]. Fransway *et.al*, 2019 reported endocrine activity, carcinogenesis, infertility, spermatogenesis, adipogenesis, perinatal exposure impact, and non-allergologic cutaneous, psychologic, and ecologic effects with parabens [13]. For this reason, many countries have banned the use of some parabens in personal care products intended for newborns and children [14,15]. FDA dissuades pregnant/nursing women and small children from using personal care products containing chlorphenesin (anti-fungal and anti-bacterial preservative) [16]. The sensitizing reactions with Amidopropyl Dimethylamines (anti-static agent) was due to the impurities like 3,3-dimethylaminopropylamine [17]. Two of the 16 main ingredients used in OTC sunblock products are safe, the FDA said. Moreover, the FDA is requesting more information on 12 ingredients among the 16 [18]. Dry skin often occurs in the elderly and tends to worsen in association with the use of standard alkaline bar soaps. Lipid solvents such as acetone, alcohols and even nonionic surfactants can cause dryness of the skin [19]. The major concern regarding the biocide triclosan (commonly used in household and personal care items to prevent the microbial growth) is antibiotic resistance [20-22]. Lee *et.al*, 2019 reported that Triclosan in toothpaste, soap, and cosmetics may accumulate in the body, which could have an adverse effect on thyroid hormones and genitalia, and increase the risk of developing breast cancer [23]. Pregnant women are particularly vulnerable to the potential risks of the endocrine disruptors contained in cosmetics [1]. Li *et.al*, 2019 reported that maternal prenatal exposure to phenols (2,4-dichlorophenol and 2,5-dichlorophenol) is inversely correlated with male birth weight and exposure to phthalates is associated with preterm birth or pregnancy loss [24].

According to Crobeddu *et.al*, 2019 phthalates family (used for industrial plasticizers to impart flexibility and durability to polyvinyl chloride) is generally considered as an endocrine disruptor could potentially increase the risks to develop breast cancer [25]. A Missouri jury ordered Johnson & Johnson to pay \$72 million to the family of Jacqueline Fox, whose death by ovarian cancer was linked to her daily use of talcum-based Johnson's Baby Powder and Shower to Shower products [26]. Park *et.al*, 2019 reported that volatile organic compounds (VOCs) and phthalates concentrations of sanitary pads and diapers were significantly higher than those found in common commercial plastic products, potentially posing risks to women (them likely causing menstrual irregularities) and children who use [27]. Diazolidinyl urea and imidazolidinyl urea are known formaldehyde releasers. The data from FDA showed that 20% cosmetic products are formaldehyde-releasers and among them, imidazolidinyl urea is the most widely used [28]. Chronic exposure to benzalkonium chloride has deleterious effects via oxidative stress, leading to cholinergic neurotoxicity [29]. Sensitization to newer and lesser known isothiazolinones has been reported [30]. Sukakul *et.al*, 2019 reported that despite the ministerial regulations restricting their use, methylchloroisothiazolinone and/or methylisothiazolinone are still found in cosmetics sold on the Thai market [31]. Studies from Europe and Israel have indicated that products with undeclared formaldehyde and formaldehyde releasers on product labels may have detectable levels of formaldehyde. For consumers allergic to formaldehyde and suffering from any kind of dermatitis, it is very important to know the potential for formaldehyde exposure in order to avoid allergic contact dermatitis (ACD) [32,33]. However, Bilal *et.al*, 2019 discussed cytotoxicity, genotoxicity, mutagenicity, neurotoxicity, estrogenicity of many of these mentioned ingredients [34]. Beauty products used for cosmetic purposes can have adverse effects to human health due to the fact that they contain Pb, Cd and other highly toxic heavy metals [35]. Cosmetics contaminated with heavy metals can lead them to accumulate in the skin, while the further migration of these toxic elements to blood vessels can cause subsequent difficulties. Saadatzadeh *et.al*, 2019 reported that arsenic contents of lipsticks, eye shadows, and eyebrow pencils was significantly higher than the BVL (Federal Office of Consumer Protection and Food Safety of Germany) standard [36]. Among the heavy metal impurities, mercury, arsenic, lead, cobalt, antimony, cadmium, nickel and chromium are highly toxic and are banned in cosmetics to be added intentionally as ingredients in EU and US [37].

The sales of skin lightening products increased 100% every year (2007-2012) in Malaysia [11]. A recent survey showed that 80% of Indian men use fairness creams and number of users growing at a rate of 20% annually. There were no differences between women and men currently using products in their desire to look as fair as media celebrities [38]. People use steroid containing pimple creams indiscriminately without knowing that steroids take off the protective outer layer of the skin so it is more exposed to UV rays and environmental pollutants such as smog and cigarette smoke [39]. In the field of dermatology, Botox A is generally injected into the muscles of facial expression for reducing the appearance of facial wrinkles [40]. Serious adverse events included dysphagia, respiratory compromise, generalized muscle weakness, marked bilateral ptosis, pseudoaneurysm of the frontal branch of the temporal artery, necrotizing fasciitis, sarcoidal granuloma, Fournier gangrene, and cervical kyphosis. Death was attributed to botulism or anaphylactic shock [41]. Applying kajal (also known as Kohl or Surma) to babies' eyes is an old tradition in many cultures of the world specially in South Asia. Most of kajal brands contains lead as one its important ingredient [42]. In people with tattoos containing red pigment of the origin of mercuric sulfur (cinnabar-vermilion, Chinese red), they may experience inflammation that is limited to this

region within 6 months of tattooing [43]. Sindoor, a cosmetic powder used in Hindu religious and cultural ceremonies has unsafe levels of lead [44].

Beauty salon workers and consumers suffered such injuries as eye and nervous system disorders, respiratory tract problems, chest pain, vomiting and rash as a result of using the straightener [45]. The percutaneous absorption of N-nitrosodiethanolamine (NDELA), an impurity in many cosmetic products, has been evaluated in diffusion cells using excised human skin [46]. p-Phenylenediamine (PPD), which was used in more than 80% of permanent dyes, is frequently reported to cause vascular neuroedema, acute renal failure or bladder cancer [47]. Qin *et.al*, 2019 reported that hair dye use increases the risk of non-Hodgkin's lymphoma, especially for females [48]. OTC use of cosmetic lenses is rapidly increasing. Several studies have reported an increased risk of infectious keratitis with the use of soft contact lenses. Prompt treatment of microbial keratitis is important to prevent vision loss [49]. Łodyga-Chruścińska *et.al*, 2018 reported presence of lead and nickel in lipsticks and powders at level prohibited by European regulation in Polish market [50]. However, several studies revealed that human exposure to heavy metals and many other potential toxins were not only from food, water, mining activity, toys, plastics, jewelry, but also from personal care products. Cosmetics are one of the most important sources of releasing heavy metals and other toxins. Different varieties of chemicals are used in cosmetic products as ingredients and some are used as preservatives. There are concerns regarding the presence of harmful chemicals in these products. Although the target hazard quotients and the hazard indices suggest a measure of safety, cosmetics may add to the body burden of potential toxic chemicals after chronic exposure.

ACKNOWLEDGEMENT

I'm Thankful to Dr. Maria L Colavincenzo, Assistant Professor of Dermatology, Department of Dermatology, Northwestern University Feinberg School of Medicine, Chicago, Illinois. For his Valuable time to Audit my Paper and for her Thoughtful Suggestions. I'm Also Grateful to Seminar Library of Faculty of Pharmacy, University of Dhaka And BANSDOC Library, Bangladesh for Providing Me Books, Journal and Newsletters.

Abbreviations

Volatile organic compounds (VOCs); allergic contact dermatitis (ACD); N-nitrosodiethanolamine (NDELA); p-Phenylenediamine (PPD)

Financial Disclosure or Funding: N/A

Conflict of Interest: The author declares that he has no competing interests.

Informed Consent: N/A

Author contributions: N/A

REFERENCES

1. Stocker, E.G. (2018). Can Makeup Harm My Vision? *Lorain NearSay*, 24(8).
2. Wang, M. T., & Craig, J. P. (2018). Investigating the effect of eye cosmetics on the tear film: current insights. *Clinical Optometry*, 10, 33.
3. Mohiuddin, A.K. Modern Age Cosmetics: An Extensive Review. *Research and Advances in Pharmacy and Life Sciences*, 1(2), 47-92
4. Marie, C., Cabut, S., Vendittelli, F., & Sauvart-Rochat, M. P. (2016). Changes in cosmetics use during pregnancy and risk perception by women. *International journal of environmental research and public health*, 13(4), 383.
5. Cornell, E. M., Janetos, T. M., & Xu, S. (2019). Time for a makeover-cosmetics regulation in the United States. *Journal of Cosmetic Dermatology*.
6. Toklu, H. Z., Antigua, A., Lewis, V., Reynolds, M. T., & Jones, J. (2019). Cosmetovigilance: A review of the current literature. *Journal of family medicine and primary care*, 8(5), 1540.
7. Mohiuddin, A.K. (2019). Cosmetics in use: a pharmacological review. *J Dermat Cosmetol*, 3(2), 50-67.
8. Mohiuddin, A. K. (2019). An Extensive Review of Sunscreen and Suntan Preparations. *OSP J Clin Trials*, 1.
9. Sautebin, L. (2008). Understanding the Adverse Effects of Cosmetics. *Drug Safety*, 31(5), 433-436.
10. Panico, A., Serio, F., Bagordo, F., Grassi, T., Idolo, A., De Giorgi, M., ... & De Donno, A. (2019). Skin safety and health prevention: an overview of chemicals in cosmetic products. *Journal of Preventive Medicine and Hygiene*, 60(1), E50.
11. Mohiuddin, A.K. (2019). Skin Lighteners & Hyperpigmentation Management. *ASIO Journal of Pharmaceutical & Herbal Medicines Research (ASIO - JPHMR)*, 5(1), 01- 42.
12. Mirza, N. (2015). Skin lightening hazards. *The Daily Star*, 8(3),
13. Fransway, A. F., Fransway, P. J., Belsito, D. V., & Yiannias, J. A. (2019). Paraben toxicology. *Dermatitis*, 30(1), 32-45.
14. Commission Regulation (EU) no. 1004/2014 of 18 September 2014 amending Annex V to Regulation (EC) no. 1223/2009 of the European Parliament and of the Council on cosmetic products. *Official Journal of the European Union*.
15. Adamson, A.(2019). Parabens: Harsh Cancer-Causing Chemicals. *Rustic Strength*, 18(3).

16. Chlorphenesin as Used in Cosmetics. Scientific Literature Review, November 28, 2011.
17. Boyer, I., Burnett, C. L., Bergfeld, W. F., Belsito, D. V., Hill, R. A., Klaassen, C. D., ... & Snyder, P. W. (2018). Safety assessment of PEGs cocamine and related ingredients as used in cosmetics. *International journal of toxicology*, 37(2_suppl), 10S-60S.
18. Mohiuddin, A.K. (2019). Sunscreen and Suntan Preparations. *ARC Journal of Pharmaceutical Sciences (AJPS)*, 5(2), 8-44.
19. Ganceviciene, R., Liakou, A. I., Theodoridis, A., Makrantonaki, E., & Zouboulis, C. C. (2012). Skin anti-aging strategies. *Dermato-endocrinology*, 4(3), 308-319.
20. Karmakar, S., Abraham, T. J., Kumar, S., Kumar, S., Shukla, S. P., Roy, U., & Kumar, K. (2019). Triclosan exposure induces varying extent of reversible antimicrobial resistance in *Aeromonas hydrophila* and *Edwardsiella tarda*. *Ecotoxicology and environmental safety*, 180, 309-316.
21. Li, M., He, Y., Sun, J., Li, J., Bai, J., & Zhang, C. (2019). Chronic Exposure to an Environmentally Relevant Triclosan Concentration Induces Persistent Triclosan Resistance but Reversible Antibiotic Tolerance in *Escherichia coli*. *Environmental science & technology*, 53(6), 3277-3286.
22. Alfihili, M. A., & Lee, M. H. (2019). Triclosan: An Update on Biochemical and Molecular Mechanisms. *Oxidative medicine and cellular longevity*, 2019.
23. Lee, J. D., Lee, J. Y., Kwack, S. J., Shin, C. Y., Jang, H. J., Kim, H. Y., ... & Kim, K. B. (2019). Risk Assessment of Triclosan, a Cosmetic Preservative. *Toxicological research*, 35(2), 137.
24. Li, H., Zheng, J., Wang, H., Huang, G., Huang, Q., Feng, N., & Xiao, J. (2019). Maternal cosmetics use during pregnancy and risks of adverse outcomes: a prospective cohort study. *Scientific reports*, 9(1), 8030.
25. Crobeddu, B., Ferraris, E., Kolasa, E., & Plante, I. (2019). Di (2-ethylhexyl) phthalate (DEHP) increases proliferation of epithelial breast cancer cells through progesterone receptor dysregulation. *Environmental research*, 173, 165-173.
26. Malkan, S. (2016). Johnson & Johnson Is Just the Tip of the Toxic Iceberg. *Time Magazine*, 2 (3).
27. Park, C. J., Barakat, R., Ulanov, A., Li, Z., Lin, P. C., Chiu, K., ... & Ko, C. J. (2019). Sanitary pads and diapers contain higher phthalate contents than those in common commercial plastic products. *Reproductive Toxicology*, 84, 114-121.
28. Ryu, O., Park, B. K., Bang, M., Cho, K. S., Lee, S. H., Gonzales, E. L. T., ... & Kim, K. B. (2018). Effects of several cosmetic preservatives on ROS-dependent apoptosis of rat neural progenitor cells. *Biomolecules & therapeutics*, 26(6), 608.
29. Antunes, S. C., Nunes, B., Rodrigues, S., Nunes, R., Fernandes, J., & Correia, A. T. (2016). Effects of chronic exposure to benzalkonium chloride in *Oncorhynchus mykiss*: cholinergic neurotoxicity, oxidative stress, peroxidative damage and genotoxicity. *Environmental toxicology and pharmacology*, 45, 115-122.
30. Herman, A., Aerts, O., de Montjoye, L., Tromme, I., Goossens, A., & Baeck, M. (2019). Isothiazolinone derivatives and allergic contact dermatitis: a review and update. *Journal of the European Academy of Dermatology and Venereology*, 33(2), 267-276.
31. Sukakul, T., Kanchanapenkul, D., Bunyavaree, M., Limphoka, P., Kumpangsin, T., & Boonchai, W. (2019). Methylchloroisothiazolinone and/or methylisothiazolinone in cosmetic products—A market survey. *Contact dermatitis*, 80(2), 110-113.
32. Nikle, A., Ericson, M., & Warshaw, E. (2019). Formaldehyde release from personal care products: chromotropic acid method analysis. *Dermatitis*, 30(1), 67-73.
33. Malinauskiene, L., Blaziene, A., Chomiciene, A., & Isaksson, M. (2015). Formaldehyde may be found in cosmetic products even when unlabelled. *Open Medicine*, 10(1).
34. Bilal, M., & Iqbal, H. M. (2019). An insight into toxicity and human-health-related adverse consequences of cosmeceuticals—a review. *Science of the total environment*.
35. Kaličanin, B., & Velimirović, D. (2016). A study of the possible harmful effects of cosmetic beauty products on human health. *Biological trace element research*, 170(2), 476-484.
36. Saadatzadeh, A., Afzalan, S., Zadehdabagh, R., Tishezan, L., Najafi, N., Seyedtabib, M., & Noori, S. M. A. (2019). Determination of heavy metals (lead, cadmium, arsenic, and mercury) in authorized and unauthorized cosmetics. *Cutaneous and ocular toxicology*, 1-5.
37. Not Just Virat Kohli, Here Are Other Celebs Who Said No To Endorsements On Ethical Grounds. *SCOOPWHOO*, Sep 14, 2017.
38. Jose, A., & Ray, J. G. (2018). Toxic content of certain commercially available fairness creams in Indian market. *Cogent Medicine*, 5(1), 1433104.
39. Mohiuddin, A.K. (2019). A Comprehensive Review of Acne Vulgaris. *Clin Res Dermatol Open Access*, 6(2), 1-3.
40. Satriyasa, B. K. (2019). Botulinum toxin (Botox) A for reducing the appearance of facial wrinkles: a literature review of clinical use and pharmacological aspect. *Clinical, cosmetic and investigational dermatology*, 12, 223.
41. Yiannakopoulou, E. (2015). Serious and long-term adverse events associated with the therapeutic and cosmetic use of botulinum toxin. *Pharmacology*, 95(1-2), 65-69.
42. Khan, F. (2019). Is surma safe for newborn eyes? *Daily Times (Pakistan)*, 24 (3),
43. Unsal, V. (2018). Natural Phytotherapeutic Antioxidants in the Treatment of Mercury Intoxication-A Review. *Advanced pharmaceutical bulletin*, 8(3), 365..
44. Shah, M. P., Shendell, D. G., Strickland, P. O., Bogden, J. D., Kemp, F. W., & Halperin, W. (2017). Lead content of sindoor, a Hindu religious powder and cosmetic: New Jersey and India, 2014–2015. *American journal of public health*, 107(10), 1630-1632.
45. Vogel, L. (2011). US legislators propose crackdown on toxic cosmetics.

46. Bronaugh, R. L., Congdon, E. R., & Scheuplein, R. J. (1981). The effect of cosmetic vehicles on the penetration of N-nitrosodiethanolamine through excised human skin. *Journal of Investigative Dermatology*, 76(2), 94-96.
47. Liu, B., Jin, S. F., Li, H. C., Sun, X. Y., Yan, S. Q., Deng, S. J., & Zhao, P. (2019). The Bio-Safety Concerns of Three Domestic Temporary Hair Dye Molecules: Fuchsin Basic, Victoria Blue B and Basic Red 2. *Molecules*, 24(9), 1744.
48. Qin, L., Deng, H. Y., Chen, S. J., & Wei, W. (2019). A Meta-Analysis on the Relationship Between Hair Dye and the Incidence of Non-Hodgkin's Lymphoma. *Medical Principles and Practice*, 28(3), 222-230.
49. Cosmetic soft contact lens associated ulcerative keratitis in southern Saudi Arabia. *Middle East Afr J Ophthalmol*.
50. Łodyga-Chruścińska, E., Sykuła, A., & Więdłocha, M. (2018). Hidden Metals in Several Brands of Lipstick and Face Powder Present on Polish Market. *Cosmetics*, 5(4), 57.